

## Activity of compounds of natural origin against *Alicyclobacillus acidoterrestris*, a common fruit juices contaminant

Agnieszka Tyfa<sup>1</sup>, Alina Kunicka-Styczyńska<sup>1</sup>, Justyna Dąbrowska<sup>2</sup>

<sup>1</sup> Institute of Fermentation Technology and Microbiology  
Lodz University of Technology 90-924, Lodz, Poland

<sup>2</sup> Institute of General Food Chemistry  
Lodz University of Technology, 90-924 Lodz, Poland

\*agnieszka.tyfa@dokt.p.lodz.pl

**Abstract:** *Fruit product industry struggles with emerging problem of microbial contamination with Alicyclobacillus acidoterrestris. This acidothermophilic, soil-borne and sporeforming bacterium posses ability to survive commercial pasteurization and thus may cause fruit juices spoilage. Even modern technologies are not effective enough to eliminate A. acidoterrestris from the industrial environments. The green consumer attitude and safety standards suggest adaptation of natural and safe solutions. This paper summarizes the bioactivity of compounds of natural origin which could serve as anti-alicyclobacilli preservation agents maintaining stability of fruit juices.*

**Keywords:** *Alicyclobacillus acidoterrestris, essential oils, bacteriocins, fruit products spoilage, antimicrobial activity.*

### Introduction

The history of *Alicyclobacillus* sp. has started in the seventies of the XX century. In 1967 an aerobic and acidophilic endospore forming microorganism had been isolated from Tokohu hot springs (Japan) and basing on morphological features both of the cells and colonies, characterized as a member of *Bacilli* class [1, 2]. Afterwards, since 1971, similar microorganisms have been isolated from differentiated habitats simultaneously overthrowing the assumption that these specific bacteria are able to survive in acidothermophilic environments only [3]. The analysis of isolates cell wall structure showed the presence of specific chemical compounds –  $\omega$ -cyclohexane fatty acids and hopanoids which are not typical components of *Bacillus* sp. cells. Moreover, the G+C content of isolates varied substantially from other bacilli, therefore, the new species, *Bacillus acidocaldarius*, has been established [3]. Further researches conducted between 1980-1984, resulted in isolation of similar bacteria from non thermal and non acidic source (soil), as well as, from aseptically packed apple juice [4, 5].

Deinhard *et al.* (1987a) investigated their mutual relationship proceeding physiological and biochemical tests for acidophilic and thermophilic *Bacillus* isolates [6]. The presence of  $\omega$ -alicyclic fatty acids in cell membranes appeared to be non specific and not limited only to *B. acidocaldarius*, therefore, a new bacterial strain, *Bacillus acidoterrestris*, was introduced [6]. The species expressed different physical and metabolic properties than *B. acidocaldarius* like ability to grow in lower temperature, lower G+C content or utilization of erythritol, sorbitol and xylitol. In addition, Deinhard *et al.* (1987b) conducted subsequent study on a new soilborne isolate containing  $\omega$ -cycloheptane fatty acids in cell membranes creating next species – *Bacillus cycloheptanicus* [7]. Finally, Wisotzkey *et al.* (1992) performed the comparative analysis of 16S rRNA for three species and revealed their high identity (around 95.0%) and affinity to the same genus with indicating low identity to other *Bacillus* sp. [8]. Therefore, they proposed the establishment of a new genus in *Bacilli* class, *Alicyclobacillus*, which gathered all known acidothermophilic bacilli with high G+C content and  $\omega$ -alicyclic fatty acids as the major membrane component of the cells. Moreover, the nomenclature of the three species has been changed as *Alicyclobacillus acidoterrestris*, *A. acidocaldarius* and *A. cycloheptanicus*.

**Table 1.** List of species and subspecies of *Alicyclobacillus* genus

Species	References
<i>A. acidiphilus</i>	[17]
<i>A. acidocaldarius</i>	[3, 8, 18, 19]
<i>A. acidocaldarius</i> subsp. <i>acidocaldarius</i>	[3, 8]
<i>A. acidocaldarius</i> subsp. <i>rittmanni</i>	[20]
<i>A. acidoterrestris</i>	[6, 8]
<i>A. aeris</i>	[21]
<i>A. cellulosityticus</i>	[9]
<i>A. contaminans</i>	[22]
<i>A. cycloheptanicus</i>	[1, 8, 7]
<i>A. dauci</i>	[10]
<i>A. disulfidooxidans</i>	[23, 24]
<i>A. fastidiosus</i>	[22]
<i>A. ferrooxydans</i>	[25]
<i>A. herbarius</i>	[26]
<i>A. hesperidum</i>	[27]
<i>A. kakegawensis</i>	[22]
<i>A. macrosporangioides</i>	[22]
<i>A. pomorum</i>	[28]
<i>A. sacchari</i>	[22]
<i>A. sendaiensis</i>	[29]
<i>A. shizuokensis</i>	[22]
<i>A. tengchongensis</i>	[30]
<i>A. tolerans</i>	[24]
<i>A. vulcanalis</i>	[31]

Until now the research on taxonomic diversity of *Alicyclobacillus* sp. has been intensively studied. Current status of *Alicyclobacillus* genus after reclassification of several species counts for 22 species and 2 subspecies (Table 1.) with the newest ones isolated from cedarwood chips and spoiled mixed juice product [9, 10]. The reclassification included an geothermal isolate from Antarctica soil (originally *Alicyclobacillus pohliae*) and clinical isolate from women blood sample (originally *Alicyclobacillus consociatus*) into *Effusibacillus pohliae* and *Effusibacillus consociatus* respectively [11-14]. However, *A. acidoterrestris* recognized as a common threat for fruit and vegetable products industry was deeply characterized [13, 15, 16].

### ***Alicyclobacillus acidoterrestris* – species characterization**

*A. acidoterrestris* is considered as a representative of *Alicyclobacillus* genus belonging to *Alicyclobacillaceae* family in *Bacillus* order. First reports on this bacterium appear in 1987, when Deinhard *et al.* isolated an unknown acidothermophilic microorganism from soil [6]. Later, a similar species was isolated from a spoiled pasteurized apple juice [5]. Current studies on morphological and physiological analyses present that *A. acidoterrestris* is an aerobic, motile, rod-shaped bacterium able to form oval endospores located centrally or terminally and may cause cell deformation [2]. The size of the endospores varies between 1.5-1.8  $\mu\text{m}$  in length and 0.9-1.0  $\mu\text{m}$  in width whereas cell size reaches 2.9-4.3  $\mu\text{m} \times 0.9$ -1.0  $\mu\text{m}$  [2, 4, 6, 32]. These bacteria remain Gram-positive, however, may stain as Gram-variable as well [7, 8].

*A. acidoterrestris* is an acidothermophilic bacterium growing in low pH and elevated temperature. The growth is observed in pH between 2.0 to 6.0 with the optimum rate at pH 4.0-5.0, however, soilborne isolates are able to survive at pH 7.0 [33, 34]. On the other hand, the optimum temperature for *A. acidoterrestris* is 40-53°C but the cells remain active in temperature range 20-70°C [4, 6, 33-35]. Major components of cellular membranes of cells and spores are  $\omega$ -alicyclic fatty acids, mainly  $\omega$ -cyclohexane C<sub>17:0</sub>, C<sub>19:0</sub>,  $\omega$ -cycloheptane, iso C<sub>15:0</sub>, C<sub>17:0</sub> and anteiso C<sub>15:0</sub>, C<sub>17:0</sub> acids [16, 29]. The genetic sequence data for 16S rRNA analysis for the microorganisms show that the overall guanine and cytosine content reaches 52.2 mol% of the genome [8, 36].

*A. acidoterrestris* is considered as a major threat for fruit processing industry. The bacterium is natural inhabitant of soil where it may survive inconvenient conditions. However, *A. acidoterrestris* might contaminate surface of vegetables and fruits especially when windfalls are collected for fruit products processing [15]. Several reports from all over the world have confirmed the presence of alicyclobacilli in acidic beverages (iced tea, juice drinks, etc.) and fruit juices made from pears, grapefruits, mango, white grapes, lemons, tomatoes and others with apple juice most often contaminated [16, 34, 37].

Juices contaminated with alicyclobacilli very rarely manifest with taint and/or precipitate, thus the spoilage cannot be detected easily [16]. The most characteristic feature of a spoiled product are specific off-flavors identified as disinfectant-like or medicinal smell [35]. This is associated with chemical metabolites produced by *Alicyclobacillus* sp., predominately guaiacol and halophenols: 2,6-DCP (2,6-dichlorophenol) and 2,6-DBC (dibromophenol) [38, 39]. The precursors of guaiacol are ferulic acid and its metabolic pathway products (vanillic acid and vanillin) as well as tyrosine [40]. Still the concentration of volatile metabolites is strictly related to a kind of juice and a number of bacterial cells [39].

The attempts to prevent *Alicyclobacillus* sp. contamination have been undertaken worldwide, however, specific acidothermophilic spores are extremely hard to eliminate. Conventional and unconventional methods involve application of techniques like high temperature (pasteurization), elevated pressure (HPH – high pressure homogenization), addition of preservation agents, ultrasounds, infrared radiation, clarification, high hydrostatic pressure (HHP) or ultrafiltration [39, 41-45]. Moreover, a promising technique authorized by U.S. Food and Drug Administration (USFDA) seems to be cleansing surface of fruits with aqueous chlorine dioxide (ClO<sub>2</sub>) and its addition to condensation water used for sanitizing technological installations [46]. On the contrary, chemicals which are approved by FDA as food additives are several enzymes, plant and bacterial metabolites which are regarded as natural compounds of confirmed antimicrobial potential. This include bacteriocins, lysozyme and plant derivatives – extracts and essential oils.

### **Bacteriocins active against *Alicyclobacillus* sp.**

Bacteriocins are considered as proteinaceous metabolites produced by Gram-positive and Gram-negative bacteria, expressing activity against microorganisms [47, 48]. Due to their high bactericidal and bacteriostatic potential, food products with bacteriocins are less susceptible to spoilage. Moreover, majority of these chemical compounds express thermostability and remain active in relatively wide pH range, therefore, are recognized as natural preservatives and could be regarded as food products additives. However, current status of bacteriocins Generally Recognized as Safe (GRAS) accounts for nisin only [47-50].

#### ***Nisin***

One of bacteriocins with well-documented inhibitory effect against Gram-positive bacteria and pathogens is nisin, a peptide produced by *Lactococcus lactis* subsp. *lactis*. Its antibacterial activity towards *A. acidoterrestris* has also been extensively studied [43, 51, 52]. Nisin is regarded as a potential preservative and suitable natural additive to fruit drinks and beverages of high acidity [53, 54]. Due to stability in both low pH and elevated temperature, nisin can be added directly into juices prior to pasteurization in order to control both *A. acidoterrestris* spores germination and growth of vegetative cells [53]. Moreover, Buonocore et al. (2004) reported that release and maintenance of nisin biological properties

could be achieved during product storage [55]. The attempts of nisin incorporation as a component of active packaging have been considered as well [56]. The amount of nisin and its effectiveness varies substantially among the studies [43, 53, 54]. These differences are caused by variety of media applied, chemical composition of juices and particular *Alicyclobacillus* sp. strain.

Nisin in concentration 100 IU/ml was proved to be effective enough to prevent growth of *A. acidoterrestris* spores and vegetative cells in apple, orange and grapefruit juices even at 45°C. Furthermore, at ambient temperature lower concentration of nisin (50 IU/ml) enhanced thermal sensitivity of *A. acidoterrestris* spores, thus prevented their outgrowth in a higher extent [51]. The 100 IU/ml dosis of nisin was also confirmed as a suitable apple and orange juice preservative in other studies [57-59].

Analogous dependence, therefore, weaker activity towards vegetative cells was observed by Yamazaki *et al.* (2000) [53]. In addition, greater acidity facilitated improvement of nisin bacteriostatic activity. Both for vegetative cells and spores cultivated on YPGA medium, the Minimum Inhibitory Concentration (MIC) values were 1.56-25 IU/ml and 25-100 IU/ml at pH 3.4 and 4.2, respectively. On the contrary, the inhibitory effect for suspension of spores only was <0.78-12.5 IU/ml at pH 3.4 and 25-100 IU/ml at pH 4.2. No outgrowth of spores was observed after 7 days of incubation for orange and mixed fruit juices with 25 IU/ml addition of nisin. However, similar effect was not achieved for the clear apple juice even with nisin dose equal to 600 IU/ml [53].

As suggested by Splittstoesser *et al.* (1994), the chemical composition of a juice itself may have an influence on microorganisms cultivation [60]. The higher the content of soluble solids and thus lower water activity ( $a_w$ ), the lower concentration of nisin is required for *A. acidoterrestris* growth inhibition [58]. Researches considering operation on several parameters (soluble solids concentration, pH, temperature) indicate that independently on the parameter, nisin at concentration 70 IU/ml is an effective inhibiting agent in both apple and orange juices of pH 3.5-5.5 and 11-19°Brix incubated at the temperature range 25-50°C [61].

On the other hand, Sokolowska *et al.* (2012) presented results of nisin effect on *A. acidoterrestris* strains isolated from polish fruit products [54]. Bacterial inhibition was observed both in 25 and 45°C with MIC value for spores germination 100-1500 IU/ml whereas for vegetative cells 50-1250 UI/ml. The results are not in agreement with study presented by Komitopoulou *et al.* (1999) [51] and Yamazaki *et al.* (2000) [53] and simultaneously indicate that polish strains of *A. acidoterrestris* are rather sensitive to higher concentration of nisin [54].

The potential bactericidal and bacteriostatic activity of nisin was corroborated in Ruiz *et al.* (2013) study considering Brazilian *A. acidoterrestris* [62]. The MIC and Minimum Bactericidal Concentration (MBC) values were assigned for 15.6 µg/ml and 31.25 µg/ml bacteriocin concentration respectively.

Moreover, the mixture of nisin and chloroform extract of *Piper aduncum* tested resulted in a synergistic effect [62].

Nisin is regarded as the bacteriocin with biological activity against *A. acidoterrestris* most intensively tested since 1999. However, apart from nisin other bacteriocins have been suggested for commercial application as biopreservative substances preventing spoilage caused by this acido-thermophilic bacteria.

### **Enterocin AS-48**

Enterocin AS-48 is a cyclic peptide extracted from *Enterococcus faecalis* A-48-32. The research by Grande *et al.* (2005) [63] confirmed antibacterial activity of enterocin against *Alicyclobacillus* sp. inoculated in fresh-made and commercial acidic juices. The concentration of bacteriocin of 2.5 µg/ml reduced the risk of *A. acidoterrestris* overgrowth in natural apple and orange juices. Spores germination in commercial juices (apple, grapefruit, orange, peach and pineapple) incubated in temperature 37°C for 90 days was inhibited and the viable cell count was reduced below the detection limit. Similarly, for enterocin-treated juices inoculated with vegetative cells at least bacteriostatic effect for 60 days of incubation was maintained. Furthermore, as suggested by Grande *et al.* (2005), addition of enterocin AS-48 to commercial acidic juices could be combined with heat treatment of lower intensity [63].

### **Warnericin RB4**

In another study Minamikawa *et al.* (2005) has presented antagonistic effect of newly characterized bacteriocin extracted from *Staphylococcus warneri* RB4 – warnericin RB4 [64]. Bacteriocin was tested for thermal and acidic stability as well as against broad microorganisms spectrum. Warnericin expressed antibacterial activity towards *Micrococcus luteus*, *A. acidoterrestris* and *A. acidocaldarius* strains. Despite strong anti-alicyclobacilli potential there are limited data on additional studies.

### **Bificin C6165**

*Bifidobacterium animalis* subsp. *animalis* CICC 6165 produces a specific IIa class bacteriocin – bificin C6165. Its antimicrobial activity against twenty strains of *Alicyclobacillus* sp. was tested [65]. The bacteriostatic effect was observed for bificin concentration at 32-64 AU/ml whereas at the concentration 512 AU/ml no viable cells were detected. The activity of bificin against alicyclobacilli was maintained only in acidic pH. Bificin thermostability up to 80°C seems to be promising in its industrial application as a biopreservative of fruit juices [65]. In the subsequent research by Pei *et al.* (2014) [66], the activity of bificin C6165 was tested against *Alicyclobacillus* sp. in diluted apple juice. Vegetative cells of *A. acidoterrestris* required at least 40 µg/ml of bificin to be reduced below the detection limit. On the contrary, for commercial fruit juices inoculated with endospores suspension, their inactivation was observed at the concentration of

80 µg/ml. The same amount of bificin added to commercial apple juice was not effective. Similar tendency, therefore lack of activity for commercial apple juice, was observed in Yamazaki *et al.* (2000) study but with nisin addition [53].

### **Paracin C**

*Lactobacillus paracasei* CICC 20241 is a particular strain in group of LAB (lactic acid bacteria) classified as a probiotic producer of bacteriocin, paracin C. In the research conducted by Pei *et al.* (2013), the biological properties as well as antimicrobial activity against *A. acidoterrestris* have been investigated [67]. Paracin C appeared to express effectiveness comparable to this of nisin, therefore, both bacteriocins are regarded as thermostable agents maintaining stability in acidic and neutral pH [51, 53, 67]. Bacteriostatic effect was observed for vegetative cell culture at the concentration 80 AU/ml whereas concentration of 160 AU/ml seemed to be the MBC value. Moreover, the amount of paracin equal to 400 AU/ml resulted in reduction of *A. acidoterrestris* endospores number by 6 log cfu/ml [67].

### **Biovicin HC5**

Biovicin HC5 is produced by *Streptococcus bovis* HC5 which expresses antimicrobial activity against many spoilage and pathogenic bacteria contaminating food products [68]. Its stability in high temperature and acidic environment has been proven, thus biovicin seems to have potential against thermoacidophilic bacteria as well [68, 69]. De Carvalho *et al.* (2008) observed that biovicin in concentration 80 AU/ml expressed bactericidal and sporicidal effect in mango pulp [68]. Moreover, biovicin increased the heat susceptibility of spores and thus the lethality of *A. acidoterrestris* cells [68].

### **Controlling enzymes – lysozymes**

Lysozymes are considered as a particular group of hydrolytic enzymes which possess an ability to 'lyse' (digest) bacterial cell wall [70]. Therefore, they might play particular role in prevention of microbial growth in food products and simultaneously enhance prolongation of food shelf stability [71, 72]. The bioactivity of lysozyme against *A. acidoterrestris* has been studied in laboratory conditions as well [55, 73-75].

Although the reports confirm its inhibition against alicyclobacilli growth, the exact mechanism seems not to be precisely specified. Bevilacqua *et al.* (2014) indicates that lysozyme could initially promote spore germination and later act on the outgrown spores causing the reduction of their amount [73]. Still the researches proceeded for both vegetative cells and spores sustain that endospores remain more sensitive to the enzyme [75]. Moreover, similarly to nisin, the exact activity of lysozyme is dose-dependent and varies within external conditions applied [73-75].

## Plant derivatives acting against alicyclobacilli

Plants are the most abundant source of natural antimicrobials, essential oils, plant extracts and hydrolates [76-78]. Essential oils are regarded as oily mixture of various chemical compounds which gained GRAS status (Generally Recognized As Safe) and approval of Food and Drug Administration (FDA) as food additives [78, 79]. They compose of several active components, however, the most potent are terpenes, alcohols, phenols, aldehydes and ketones [79]. Apart from essential oils, their by-products are collected during the production as well. These includes hydrolates and plant extracts, however, the oils themselves are applied mostly.

Essential oils are reported to exhibit biological activity against microorganisms. This might be expressed either by microbial growth inhibition or as elimination and cidal effect [77, 79]. Fruit juices are commonly spoiled by *Alicyclobacillus* bacteria, however, several researches have considered application of essential oils or their active constituents as natural antimicrobial agents [52, 80-82]. According to Maldonado *et al.* (2013), lemon essential oil seems to be potential inhibitor of alicyclobacilli even up to 11 days of incubation in culture media [81]. However, the efficiency of the oil could depend on its exact chemical composition and production method [81]. On the contrary, Huertas *et al.* (2014) [52] and Bevilacqua *et al.* (2010) [82] conducted the research involving active components of essential oils only. According to study by Huertas *et al.* (2014) on citrus essential oils components, only 0.69 mM of citral could decrease the amount of *A. acidoterrestris* spores by 1 log cfu/ml, whereas limonene even at concentration 3.7 mM appeared to be ineffective. However, combining two natural antimicrobials citral and nisin (1.5 mg/L and 0.69 mM respectively) with heating in 95°C for 2 minutes led to complete elimination of the bacteria [52]. Research by Bevilacqua *et al.* (2010) considered the most active components of cinnamon and clove essential oils – eugenol and cinnamaldehyde, added into the apple juice [82]. The preservative effect was achieved for this mixture at concentration 40 ppm of eugenol and 20 ppm of cinnamaldehyde, however, the better stability of juice was observed for their doubled concentrations [82].

Apart from essential oils, plant extracts were tested for their antimicrobial potential as well [83,84]. Bevilacqua *et al.* (2013) compared the bioactivity of citrus extracts on *A. acidoterrestris* spores number during thermal processing [83]. The reduction of spores by 5 log cfu/ml was observed for all three extracts tested (lemon, neroli, bicitro) at concentration 500 ppm. Bioactivity of these extracts seemed to be strain dependent, however, higher concentrations applied should also suppress the growth of any additional microbiota. The MIC values achieved for bicitro have fallen in the range 250-500 ppm, whereas for lemon extracts lower concentration was required to maintain bacteriostatic effect (160-250 ppm). Neroli seemed, however, inadequate as an apple juice additive due to substantial alternation of juice organoleptic features [83].



Molva and Baysal (2015) investigated the effect of grape seed extract on *A. acidoterrestris* vegetative cells and spore viability [84]. The addition of 0.23-3.6% extract to apple juice stored in 37°C for 14 days resulted in the decrease of microbial population by 3-4 log cfu/ml, which seems to be promising [84].

An interesting research was proceeded by Alberice *et al.* (2012) and involved usage of chemically active glycosides saponins [85]. The inactivation of alicyclobacilli was tested for both commercial saponin and purified methanolic extracts from *S. saponaria* and obtained results seemed comparable. Treatment with 100 mg/L commercial saponin reduced the amount of spores by 2.34 log, whereas 500 mg/L purified plant extract maintained juice stability longer than for 5 days. According to these studies, saponins may be also used in preservation of fruit juices.

## Summary

Presence of *A. acidoterrestris* is an emerging problem for the fruit products industry. This acidothermophilic and sporeforming microorganisms is able to spoil pasteurized juices causing them inadequate for the consumption. Therefore, prevention and elimination methods are of a very high concern. Current technologies involving thermal treatment, adaptation of high pressure, sanitizing of industrial installations are not effective enough. The best solution seems to be direct action oriented towards *A. acidoterrestris* cells and spores by chemical agents. However, incorporation of chemical substances into food products should be strictly controlled and stay in agreement with FDA standards. Current trends for green consumption requires rejection of synthetic preservative substances and thus natural plant substances are appreciated. Nevertheless, fruit juices industry should consider only these substances which will not only maintain microbiological shelf stability of products but also will not affect product organoleptic features. Essential oils and extracts as well as bacteriocins are compounds of natural origin with proved antimicrobial activity. Moreover, majority of them had gained GRAS status and may be added directly into the food. Bacteriocins and plant metabolites effectiveness against *A. acidoterrestris* has been intensively studied since decades. According to current knowledge, the addition of any of these compounds might at least reduce the viability of the bacteria and sustain the quality and product shelf stability. Still the activity of natural plant and microbial substances remain an open topic for further researches.

## References

1. Uchino F, Doi S. Acido-thermophilic bacteria from thermal waters. *Agr Biol Chem Tokyo* **1967**, 31:817-822.
2. Yamazaki K, Teduka H, Shinano S. Isolation and identification of *Alicyclobacillus acidoterrestris* from acidic beverages. *Biosci Biotech Biochem* **1996**, 60: 543-545.
3. Darland G, Brock TD. *Bacillus acidocaldarius* sp. nov., an acidophilic, thermophilic spore-forming bacterium. *J Gen Microbiol* **1971**, 67:9-15.
4. Hippchen B, Roll A, Poralla K. Occurrence in soil of thermo-acidophilic bacilli possessing  $\omega$ -cyclohexane fatty acids and hopanoids. *Arch Microbiol* **1981**, 129:53-55.

5. Cerny G, Hennlich W, Poralla K. Fruchtsaftverderb durch *Bacillen*: isolierung und charakterisierung des verderbserregers. *Z Lebensm Unters For* **1984**, 179:224-227.
6. Deinhard G, Blanz P, Poralla K, Altan E. *Bacillus acidoterrestris* sp. nov. a new thermotolerant acidophile isolated from different soils. *Syst Appl Microbiol* **1987**, 10:47-53.
7. Deinhard G, Saar J, Krischke W, Poralla K. *Bacillus cycloheptanicus* sp. nov., a new thermoacidophile containing  $\omega$ -cycloheptane fatty acids. *Syst Appl Microbiol* **1987**, 10:68-73.
8. Wisotzkey JD, Jurtshuk P, Fox GE, Deinhard G, Poralla K. Comparative sequence analyses on the 16S rRNA (rDNA) of *Bacillus acidocaldarius*, *Bacillus acidoterrestris*, and *Bacillus cycloheptanicus* and proposal for creation of a new genus, *Alicyclobacillus* gen. nov. *Int J Syst Evol Microbiol* **1992**, 42:263-269.
9. Kusube M, Sugihara A, Moriwaki Y, Ueoka T, Shimane Y, Minegishi H. *Alicyclobacillus cellulosityticus* sp. nov., a thermophilic, cellulolytic bacterium isolated from steamed Japanese cedar chips in lumbermill. *Int J Syst Evol Microbiol* **2014**, 64:2257-2263.
10. Nakano C, Takahashi N, Okada S. *Alicyclobacillus dauci* sp. nov., a novel slightly thermophilic acidophilic bacterium isolated from a spoiled mixed vegetable and fruit juice product. *Int J Syst Evol Microbiol* **2014**, 65:716-722.
11. Watanabe M, Kojima H, Fukui M. Proposal of *Effusibacillus lacus* gen. nov., sp. nov., and reclassification of *Alicyclobacillus pohliae* as *Effusibacillus pohliae* comb. nov. and *Alicyclobacillus consociatus* as *Effusibacillus consociatus* comb. nov. *Int J Evol Microbiol* **2014**, 64:2770-2774.
12. Imperio T, Viti C, Marri L. *Alicyclobacillus pohliae* sp. nov., a thermophilic, endospore-forming bacterium isolated from geothermal soil of the north-west slope of Mount Melbourne (Antarctica). *Int J Syst Evol Microbiol* **2008**, 58:221-225.
13. Walker M, Phillips CA. *Alicyclobacillus acidoterrestris*: an increasing threat to the fruit juice industry? *Int J Food Sci Technol* **2008**, 43:250-260.
14. Glaeser SP, Falsen E, Martin K, Kampfer P. *Alicyclobacillus consociatus* sp. nov., isolated from a human clinical specimen. *Int J Syst Evol Microbiol* **2013**, 63:3623-3627.
15. Satora P. *Alicyclobacillus acidoterrestris* – przetrwalnikujaca bakteria skazajaca soki. *Laboratorium* **2009**, 5:20-24.
16. Chmal-Fudali E, Papiewska A. The possibility of thermal inactivation of *Alicyclobacillus acidoterrestris* spores in fruit and vegetable juices. *Biotechnol Food Sci* **2011**, 75:87-96.
17. Matsubara H, Goto K, Matsumara T, Mochida K, Iwaki M, Niwa M, Yamasato K. *Alicyclobacillus acidophilus* sp. nov., a novel thermoacidophilic  $\omega$ -alicyclic fatty acid-containing bacterium isolated from acidic beverages. *Int J Syst Evol Microbiol* **2002**, 52:1681-1685.
18. De Rosa M, Gambacorta A, Minale L, Bu'Lock JD. Cyclohexane fatty acids from a thermophilic bacterium. *J Chem Soc Chem Comm* **1971**, p. 1334.
19. Loginova LG, Khraptsova GI, Egorova LA, Bogdanova TI. Acidophilic, obligate thermophilic bacterium *Bacillus acidocaldarius* isolated from hot springs and soil of Kusnashir island. *Microbiology* **1978**, 47:771-775.
20. Nicolaus B, Improta R, Manca MC, Lama L, Esposito E, Gambacorta A. *Alicyclobacilli* from an unexpected geothermal soil in Antarctic: Mount Rittmann Polar Biol **1998**, 19:133-141.
21. Guo X, You XY, Liu LJ, Zhang JY, Liu SJ, Jiang CY. *Alicyclobacillus aeris* sp. nov., a novel ferrous-and sulfur-oxidizing bacterium isolated from a copper mine. *Int J Syst Evol Microbiol* **2009**, 59:2415-2410.

22. Goto K, Mochida K, Kato Y, Asahara M, Fujita R, An SY, Kasai H, Yokota A. Proposal of six species of moderately thermophilic, acidophilic, endospore-forming bacteria: *Alicyclobacillus contaminans* sp. nov., *Alicyclobacillus fastidious* sp. nov., *Alicyclobacillus kakegawensis* sp. nov., *Alicyclobacillus macrosporangioides* sp. nov., *Alicyclobacillus sacchari* sp. nov., and *Alicyclobacillus shizuokensis* sp. nov. *Int J Syst Evol Microbiol* **2007**, 57:1276-1285.
23. Dufresne S, Bousquet J, Boissinot M, Guay R. *Sulfobacillus sidulfidooxidans* sp. nov., a new acidophilic, disulfide-oxidizing, gram-positive spore-forming bacterium. *Int J Syst Evol Microbiol* **1996**, 46:1056-1064.
24. Karavaiko GI, Bogdanova TI, Tourova TP, Kondrat'eva TF, Tsaplina IA, Egorova MA, Krasil'nikova EN, Zakharchuk LM. Reclassification of '*Sulfobacillus thermosulfidooxidans* subsp. *thermotolerans*' strain K1 as *Alicyclobacillus disulfidooxidans* comb. nov., and emended description of the genus *Alicyclobacillus*. *Int J Syst Evol Microbiol* **2005**, 55:941-947.
25. Jiang CY, Liu YY, You XY, Guo X, Liu SJ. *Alicyclobacillus ferrooxydans* sp. nov., a ferrous-oxidizing bacterium from solfataric soil. *Int J Syst Evol Microbiol* **2008**, 58:2898-2903.
26. Goto K, Matsubara H, Mochida K, Matsubara T, Hara Y, Niwa M, Yamasato K. *Alicyclobacillus herbarius* sp. nov., a novel bacterium containing  $\omega$ -cycloheptane fatty acids, isolated from herbal tea. *Int J Syst Evol Microbiol* **2002**, 52:109-113.
27. Albuquerque L, Rainey FA, Chung AP, Sunna A, Nobre MF, Grote R, Antranikian G, Da Costa MS. *Alicyclobacillus hesperidum* sp. nov., and related genomic species from solfataric soils of São Miguel in the Azores. *Int J Syst Evol Microbiol* **2000**, 50:451-457.
28. Goto K, Mochida K, Asahara M, Suzuki M, Kasai H, Yokota A. *Alicyclobacillus pomorum* sp. nov., a novel thermo-acidophilic, endosporeforming bacterium that does not possess  $\omega$ -alicyclic fatty acids, and emended description of the genus *Alicyclobacillus*. *Int J Syst Evol Microbiol* **2003**, 53:1537-1544.
29. Tsuruoka N, Isono Y, Shida O, Hemmi H, Nakayama T, Nishino T. *Alicyclobacillus sendaiensis* sp. nov., a novel acidophilic, slightly thermophilic species isolated from soil in Sendai, Japan. *Int J Syst Evol Microbiol* **2003**, 53:1081-1084.
30. Kim MG, Lee JC, Park DJ, Li WJ, Kim CJ. *Alicyclobacillus tengchongensis* sp. nov., a thermo-acidophilic bacterium isolated from hot springs. *J Microbiol* **2014**, 52:884-889.
31. Simbahan J, Drijber R, Blum P. *Alicyclobacillus vulcanalis* sp. nov., a thermophilic, acidophilic bacterium isolated from Coso Hot Springs, California, USA. *Int J Syst Evol Microbiol* **2004**, 54:1703-1707.
32. Oteiza JM. *Imagenes Microbiológicas. Alicyclobacillus acidoterrestris*. *Rev Arg Microbiol* **2011**, 43:67.
33. Bevilacqua A, Corbo MR, D'Amato D, Campaniello D, Sinigaglia M. Caratterizzazione fenotipica di ceppi di *Alicyclobacillus* spp. isolati da suolo. Ricerche e innovazioni nell'industria alimentare, vol. VII. Proceedings of the VI Italian Conference of Food Science and Technology (Cernobbio-CO, Italy-19-20 September **2005**) Chiriotti Editori, Pinerolo, TO, Italy, pp. 1201-1205.
34. Sokolowska B, Niezgoda J, Chotkiewicz M. Opportunities to germinate and grow of *Alicyclobacillus acidoterrestris* spores in the presence of organic acids. *FMFI* **2013**, 2:10-16.
35. Walls I, Chuyate R. *Alicyclobacillus* – historical perspective and preliminary characterization study. *Dairy Food Environ Sanit* **1998**, 18:499-503.

36. Shemesh M, Pasvolsky R, Sela N, Green SJ, Zakin V. Draft genome of *Alicyclobacillus acidoterrestris* strain ATCC 49025. *Genome Announc* **2013**, 1(5):e00638-13.
37. Smit Y, Cameron M, Venter P, Witthuhn RC. *Alicyclobacillus* spoilage and isolation – a review. *Food Microbiol* **2011**, 28:331-349.
38. Borlinghaus A, Engel R. *Alicyclobacillus* incidences in commercial apple juice concentrate (AJC) supplies – methods development and validation. *Fruit Process* **1997**, 7:262.
39. Kumar R, Bawa AS, Kathiravan T, Nadanasabapathi S. Inactivation of *Alicyclobacillus acidoterrestris* by non thermal processing technologies – a review. *Internat J Adv Res* **2013**, 1:386-395.
40. Witthuhn RC, van der Merwe E, Venter P, Cameron M. Guaiacol production from ferulic acid, vanillin and vanillic acid by *Alicyclobacillus acidoterrestris*. *Int J Food Prot* **2012**, 157:113-117.
41. Lee SY, Dougherty RH, Kang DH. Inhibitory effects of high pressure and heat on *Alicyclobacillus acidoterrestris* spores in apple juice. *Appl Environ Microbiol* **2002**, 68:4158-4161.
42. Bahceci KS, Gokmen V, Serpen A, Acar J. The effects of different technologies on *Alicyclobacillus acidoterrestris* during apple juice production. *Eur Food Res Technol* **2003**, 217:249-252.
43. Bevilacqua A, Sinigaglia M, Corbo MR. *Alicyclobacillus acidoterrestris*: New methods for inhibiting spore germination. *Int J Food Microbiol* **2008**, 125:103-110.
44. Djas M, Bober M, Henczka M. New methods for inactivation of *Alicyclobacillus acidoterrestris* spores in apple juice concentrate. *Chall Modern Technol* **2011**, 2:4649.
45. Evrendilek GA, Baysal T, Icier F, Yildiz H, Demirdoven A, Bozkurt H. Processing of fruits and fruit juices by novel electrotechnologies. *Food Eng Rev* **2012**, 4:68-87.
46. FDA, USDA, CDC. Guidance for industry – guide to minimize microbial food safety hazards for fresh fruits and vegetables. **1998**.  
<http://www.fda.gov/downloads/Food/GuidanceRegulation/UCM169112.pdf>
47. Gwiazdowska D, Trojanowska K. Bakteriocynty – właściwości i aktywność przeciwdrobnoustrojowa. *Biotechnologia* **2015**, 68:114-130.
48. Karpiński TM, Szkaradkiewicz AK. Characteristic of bacteriocins and their application. *Pol J Microbiol* **2013**, 62:223-235.
49. FDA, CFSAN. GRAS notice information **2013**.  
<http://www.fda.gov/downloads/food/ingredientspackaginglabeling/gras/noticeinventory/ucm266587.pdf>
50. Yang SC, Lin CH, Sung CT, Fang JY. Antimicrobial activities of bacteriocins: application in foods and pharmaceuticals. *Front Microbiol* **2014**, 5:241.
51. Komitopoulou E, Boziaris IS, Davies EA, Delves-Broughton J, Adams MR. *Alicyclobacillus acidoterrestris* in fruit juices and its control by nisin. *Int J Food Sci Technol* **1999**, 34:81-85.
52. Huertas JP, Esteban MD, Antolines V, Palop A. Combined effect of natural antimicrobials and thermal treatments on *Alicyclobacillus acidoterrestris* spores. *Food Control* **2014**, 35:73-78.
53. Yamazaki K, Murakami M, Kawai Y, Inoue N, Matsuda T. Use of nisin for inhibition of *Alicyclobacillus acidoterrestris* in acidic drinks. *Food Microbiol* **2000**, 17:315-320.

54. Sokolowska B, Niezgoda J, Chotkiewicz M. Wpływ nizyny i lizozymu na wzrost szczepow *Alicyclobacillus acidoterrestris* oraz mozliwosc zastosowania tych zwiazkow jako biokonserwantow w soku jablkowym. ZYWNOSC. Nauka. Technologia. Jakosc. **2012**, 4:44-54.
55. Buonocore GG, Sinigaglia M, Corbo MR, Bevilacqua A, La Notte E, Del Nobile MA. Controlled release of antimicrobial compounds from highly swellable polymers. J Food Prot **2004**, 67:1190-1194.
56. Barbosa AA, Silva de Araujo HG, Matos PN, Carnelossi MA, Almeida de Castro A. Effects of nisin-incorporated films on the microbiological and physicochemical quality of minimally processes mangoes. Int J Food Microbiol **2013**, 164:135-40.
57. Peña WE, de Massaguer PR. Microbial modeling of *Alicyclobacillus acidoterrestris* CRA 7152 growth in orange juice with nisin addition. J Food Protect **2006**, 69:1904-1912.
58. Peña WE, de Massaguer PR, Teixeira LQ. Microbial modeling of thermal resistance of *Alicyclobacillus acidoterrestris* CRA 7152 spores in concentrated orange juice with nisin addition. Braz J Microbiol **2009**, 40:601-611.
59. Peña WE, de Massaguer PR, Zuniga AD, Saraiva SH. Modeling the growth limit of *Alicyclobacillus acidoterrestris* CRA7152 in apple juice: effect of pH, Brix, temperature and nisin concentration. J Food Process Pres **2011**, 35:509-517.
60. Splittstoesser DF, Churey JJ, Lee CY. Growth characteristics of aciduric sporeforming bacilli isolated from fruit juices. J Food Prot **1994**, 57:1080-1083.
61. Peña WE, de Massaguer PR. Modeling the growth/no growth interface of *Alicyclobacillus acidoterrestris* CRA 7152 in orange juice as a function of pH, temperature, Brix and nisin concentration. Scientia Agropecuaria **2010**, 1:47-61.
62. Ruiz SP, dos Anjos MM, Carrara VS, deLima JN, Cortez DA, Nakamura TU, Nakamura CV, de Abreu Filho BA. Evaluation of the antibacterial activity of *Piperaceae* extracts and nisin on *Alicyclobacillus acidoterrestris*. J Food Sci **2013**, 78:1772-1777.
63. Grande MJ, Lucas R, Abriouel H, Omar NB, Maqueda M, Martinez-Bueno M, Martinez-Canamero M, Valdivia E, Galvez A. Control of *Alicyclobacillus acidoterrestris* in fruit juices by enterocin AS-48. Int J Food Microbiol **2005**, 104:289-297.
64. Miamikawa M, Kawai Y, Inoue N, Yamazaki K. Purification and characterization of warnericin RB4, anti-*Alicyclobacillus* bacteriocin, produced by *Staphylococcus warneri* RB4. Curr Microbiol **2005**, 51:22-26.
65. Pei J, Yuan Y, Yue T. Characterization of bacteriocin bificin c6165: a novel bacteriocin. J Appl Microbiol **2013**, 114:1273-1284.
66. Pei J, Yue T, Yuan Y. Control of *Alicyclobacillus acidoterrestris* in fruit juices by a newly discovered bacteriocin. World J Microbiol Biotechnol **2014**, 30:855-863.
67. Pei J, Yuan Y, Yue T. Primary characterization of bacteriocin paracin C – a novel bacteriocin produced by *Lactobacillus paracasei*. Food Control **2013**, 34:168-176.
68. De Carvalho AA, Vanetti MC, Mantovani HC. Biovicin HC5 reduces thermal resistance of *Alicyclobacillus acidoterrestris* in acidic mango pulp. J Appl Microbiol **2008**, 104:1685-1691.
69. Mantovani HC, Hu H, Worobo RW, Russell JB. Biovicin HC5, a bacteriocin from *Streptococcus bovis* HC5. Microbiol **2002**, 148:3347-3352.
70. Nerurkar LS. 67-Lysozyme. In: Methods for studying mononuclear phagocytes. Adams DO, Edelson PJ, Koren HS; Academic Press INC., **1981**, pp. 667-683.

71. Proctor VA, Cunningham FE. The chemistry of lysozyme and its use as a food preservative and a pharmaceutical. *Crit Rev Food Sci Nutr* **1988**, 26:359-395.
72. Cunningham FE, Proctor VA, Goetsch SJ. Egg-white lysozyme as a food preservative: an overview. *World Poultry Sci J* **1991**, 47:141-163.
73. Bevilacqua A, Ciuffreda E, Sinigaglia M, Corbo MR. Effects of lysozyme on *Alicyclobacillus acidoterrestris* under laboratory conditions. *Int J Food Sci Technol* **2014**, 49:224-229.
74. Sokolowska B, Skapska S, Fonberg-Broczek M, Niezgoda J, Chotkiewicz M, Dekowska A, Rzoska S. The combined effect of high pressure and nisin or lysozyme on the inactivation of *Alicyclobacillus acidoterrestris* spores in apple juice. *High Pressure Res* **2012**, 32:119-127.
75. Conte A, Sinigaglia M, Del Nobile MA. Antimicrobial effectiveness of lysozyme immobilized on polyvinylalcohol-based film against *Alicyclobacillus acidoterrestris*. *J Food Prot* **2006**, 69:861-865.
76. Rabha B, Gopalakrishnan R, Baruah I, Singh L. Larvicidal activity of some essential oil hydrolates against dengue and filiriasis vectors. *EJMR* **2012**, 1:014-016.
77. Tajkarimi MM, Ibrahim SA, Cliver DO. Antimicrobial herb and spice compounds in food. *Food Control* **2010**, 21:1199-1218.
78. Burt S. Essential oils: their antimicrobial properties and potential applications in foods – a review. *Int J Food Microbiol* **2004**, 94:223-253.
79. Bevilacqua A, Corbo MR, Campaniello D, D'Amato D, Gallo M, Speranza B, Sinigaglia M. Shelf life prolongation of fruit juices through essential oils and homogenization: a review. In: Science against microbial pathogens: communicating current research and technological advances. Mendez-Vilas A.; Formatex Research Center, Badajoz, Spain, **2011**, Vol. 2, pp. 1157-1166.
80. Bevilacqua A, Corbo MR, Sinigaglia M. Use of essential oils to inhibit *Alicyclobacillus acidoterrestris*: a short overview of the literature. *Front Microbiol* **2011**, 2:195, doi:10.3389/fmicb.2011.00195.
81. Maldonado MC, Aban MP, Navarro AR. Chemicals and lemon essential oil effect on *Alicyclobacillus acidoterrestris* viability. *Braz J Microbiol* **2013**, 44:1133-1137.
82. Bevilacqua A, Corbo MR, Sinigaglia M. Combining eugenol and cinnamaldehyde to control the growth of *Alicyclobacillus acidoterrestris*. *Food Control* **2010**, 21:172-177.
83. Bevilacqua A, Campaniello D, Speranza B, Sinigaglia M, Corbo MR. Control of *Alicyclobacillus acidoterrestris* in apple juice by citrus extracts and a mild heat-treatment. *Food Control* **2013**, 31:553-559.
84. Molva C, Baysal AH. Antimicrobial activity of grape seed extract on *Alicyclobacillus acidoterrestris* DSM 3922 vegetative cells and spores in apple juice. *LWT-Food Sci Technol* **2015**, 60:238-245.
85. Alberice JV, Funes-Huacca ME, Guterres SB, Carrilho E. Inactivation of *Alicyclobacillus acidoterrestris* in orange juice by saponin extracts combined with heat-treatment. *Int J Food Microbiol* **2012**, 159:130-135.